

We claim:

1. An isolated or purified polynucleotide comprising a nucleotide sequence which encodes a human endosulfine and fragments or compliments thereof.

2. The polynucleotide of Claim 1 wherein said nucleotide is SEQ ID NO:1.

3. The polynucleotide of Claim 1 wherein said nucleotide is SEQ ID NO:2.

4. The polynucleotide of Claim 1 wherein said nucleotide is SEQ ID NO:1 from about nucleotide position 107 to about nucleotide position 460.

5. The polynucleotide of Claim 1 wherein said nucleotide is SEQ ID NO:2 from about nucleotide position 107 to about nucleotide position 472.

6. The human endosulfine of Claim 1 wherein said nucleotide sequence encodes an amino acid sequence SEQ ID NO:3 or fragments thereof.

7. The human endosulfine of Claim 1 wherein said nucleotide sequence encodes an amino acid sequence SEQ ID NO:4 or fragments thereof.

8. A recombinant expression vector comprising the polynucleotide of Claim 1.

9. A recombinant expression vector comprising the polynucleotide of Claim 4.

10. A recombinant expression vector comprising the polynucleotide of Claim 5.

11. The expression vector of Claim 8 selected from the group consisting of pProEx1 and pcDNA3.1.

12. The expression vector of Claim 9 selected from the group consisting of pProEx1 and pcDNA3.1.

13. The expression vector of Claim 10 selected from the group consisting of pProEx1 and pcDNA3.1.

buta1

10

000241380400001

20

25

30

35

buta2

14. A host cell transformed with the expression vector of Claim 8.

15. The host cell of Claim 14 wherein said host cell is a prokaryotic cell or eukaryotic cell.

16. A host cell transformed with the expression vector of Claim 9.

17. The host cell of Claim 16 wherein said host cell is a prokaryotic cell or eukaryotic cell.

18. A host cell transformed with the expression vector of Claim 10.

19. The host cell of Claim 18 wherein said host cell is a prokaryotic cell or eukaryotic cell.

20. An isolated or purified polypeptide of human endosulfine.

21. The polypeptide of Claim 20 having the sequence of SEQ ID NO:3 or fragments thereof.

22. The polypeptide of Claim 20 having the sequence of SEQ ID NO:4 or fragments thereof.

23. A method for producing a polypeptide containing at least one human endosulfine epitope comprising incubating host cells transformed with an expression vector wherein said expression vector comprises a nucleotide sequence which encodes a human endosulfine.

24. The method of Claim 23 wherein said nucleotide sequence which encodes a human endosulfine has the sequence SEQ ID NO:2 and fragments and compliments thereof.

25. The method of Claim 24 wherein said nucleotide sequence which encodes a human endosulfine has the sequence SEQ ID NO:2 from about nucleotide position 107 to about nucleotide position 472.

26. The method of Claim 23 wherein said nucleotide sequence encodes a human endosulfine having the sequence SEQ ID NO:3.

27. The method of Claim 23 wherein said nucleotide sequence encodes a human endosulfine having the sequence SEQ ID NO:4.

5
28. The method of Claim 23 wherein said nucleotide sequence which encodes a human endosulfine has the sequence SEQ ID NO:2.

10
29. The method of Claim 23 wherein said nucleotide sequence which encodes a human endosulfine has the sequence SEQ ID NO:2 from about nucleotide position 107 to about nucleotide position 472.

15
30. A method for identifying compounds that modulate endosulfine receptor activity, comprising:

- 20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995
- (a) providing a host cell that expresses said endosulfine receptor polypeptide;
 - (b) mixing a test compound with said cell; and
 - (c) measuring either
 - (i) the effect of the test compound on the cell expressing the receptor, or
 - (ii) the binding of the test compound to the cell or to the receptor.

31. The method of claim 30, wherein said host cell is either a prokaryotic or eukaryotic cell.

32. The method of claim 30, wherein said measurement of step (c) (ii) is performed by measuring a signal generated by a signal generating compound.

33. The method of claim 30, wherein the measurement of step (c) (ii) is performed by measuring a signal generated by a radiolabeled ion, a fluorescent probe or an electrical current.

34. A method for identifying a cytoprotective compound, comprising:

- (a) providing a cell that expresses an endosulfine polypeptide or fragment thereof;
- (b) combining a test compound with the cell; and
- (c) monitoring the cell or cellular function for an indication of cytotoxicity.

35. The method of claim 34 wherein said cell is either a prokaryotic or eukaryotic cell.

36. The method of claim 35, wherein said cell comprises an expression vector comprising the polynucleotide of claim 5 operably linked to control sequences that direct the transcription of the polynucleotide whereby the polynucleotide is expressed in a host cell.

5 37. The method of claim 36, wherein at least one of the control sequences comprises an inducible promotor.

10 38. The method of claim 37, wherein said cell is maintained in the presence of a substance such as to minimize or block a cytotoxic effect on said cell.

15 39. A method of treating an individual having a condition associated with endosulfine modulation, comprising administering to said individual an effective amount of a compound that controls the gene expression of endosulfine, in a pharmaceutically acceptable excipient.

40. A monoclonal antibody which specifically binds to human endosulfine having amino acid sequence SEQ ID NO:4 or fragments thereof.

Add 4

0924178-0402031
1022040-8242350